


TOM. 6 / 2015

ISSN 1804-2732



# Acta Carpathica Occidentalis

PŘÍRODA ZÁPADNÍCH KARPAT



Muzeum regionu Valašsko, Vsetín  
Muzeum jihovýchodní Moravy ve Zlíně



**First record of *Strigamia pusilla* from the Czech Republic  
(*Chilopoda: Geophilomorpha*)**  
**První nález zemivky *Strigamia pusilla* z území České republiky  
(*Chilopoda: Geophilomorpha*)**

Ivan Hadrián Tuf<sup>1</sup> & Jiří Kupka<sup>2</sup>

<sup>1</sup> Department of Ecology and Environmental Sciences, Faculty of Science, Palacký University Olomouc; e-mail: ivan.tuf@upol.cz  
<sup>2</sup> Institute of Environmental Engineering, Faculty of Mining and Geology, Technical University of Ostrava; e-mail: jiri.kupka@vsb.cz

**Keywords:** Beskydy, centipede, Linotaeniidae, Moravian-Silesian Beskids

**Abstrakt:** Zemivka *Strigamia pusilla* (Sseliwanoff, 1884) byla nalezena poprvé na území ČR, konkrétně v NPR Mazák (CHKO Beskydy). Jedná se o čtvrtý druh daného rodu známého z Česka. Její rozšíření a klíčové znaky jsou uvedeny.

**Abstract:** The centipede *Strigamia pusilla* (Sseliwanoff, 1884) was found for the first time in the Czech Republic in the Mazák National Nature Reserve, the Beskydy Protected Landscape Area. It is the fourth species of the genus *Strigamia* known from the territory of the Czech Republic. Information about its distribution and key characters are added.

## INTRODUCTION

Centipede genus *Strigamia* Gray, 1843 is rather a complicated taxon from the family Linotaeniidae (Chilopoda, Geophilomorpha), which needs taxonomical revision. Its distribution is Holarctic. Altogether, 44 valid species are recognized recently (BONATO et al. 2012), among them 12 species are distributed in Europe (BONATO & MINELLI 2013). Nevertheless, another 31 species and subspecies announced in past from Europe are synonymised with valid European species. Moreover, half of those dozen valid species were not described properly according to modern demands on the description of geophilomorph centipedes, so their validity is uncertain because their morphology is inadequately known (BONATO et al. 2010).

There were three species of the genus *Strigamia* known from the Czech Republic (TUF & TUFOVÁ 2008), *Strigamia acuminata* (Leach, 1815), *Strigamia crassipes* (C. L. Koch, 1835) and *Strigamia transsilvanica* (Verhoeff, 1928). All these three species are widely distributed

through Bohemia and Moravia (TUF & LAŠKA 2005). This paper reports the first record of another species of this genus from the territory of the Czech Republic.

## MATERIAL

*Strigamia pusilla* (Sseliwanoff, 1884): 1 male (33 pairs of legs) and 1 female (35 pairs of legs) – the Czech Republic, Moravia, Beskydy Protected Landscape Area, Mazák National Nature Reserve, debris of wood (*Tilio-Acerion* plant association) in NW part of the reserve, N:49°32'32,985", E:18°26'19,182", faunistic square code 6476D, 10.ix.2012, Jiří Kupka leg.

## DISCUSSION


All species of the genus *Strigamia* occurring on the territory of the Czech Republic are recognised as valid and certain. Distribution of *S. acuminata* and *S. crassipes* is throughout the whole Europe to the Caucasus. Distribution of *S. transsilvanica* is connected with the Carpathians, but extends to the entire Alps, to the Baltic, and the mainland of Greece respe-

Fig. 1: Plant communities of debris woods are unique due to its specific soil conditions as well as specific microclimate. Orig. Jiří Kupka.  
Obr. 1: Vegetace suťových lesů je unikátní jak díky specifickým vlastnostem, tak kvůli odlišnému mikroklimatu. Orig. Jiří Kupka.

ctively. Distribution of *S. pusilla* is going from the Sudeten through the Carpathians to Siberia and Central Asia. It was reported from Slovakia, Poland, Romania, Russia and Mongolia yet. Recently it was redescribed and in detail illustrated by DÁNYI (2006).

It is rather common to found more species of this genus together on the same locality. Also the Mazák National Nature Reserve is occupied by at least three species of the *Strigamia*, namely *S. crassipes* (1 male), and *S. acuminata* (1 female, 3 juv.) were collected here during the same sampling occasion as well. The investigated part of forest is one of the oldest growths on territory of the given Nature Reserve (Fig. 1).

Centipedes of the genus *Strigamia* are well recognizable among our centipedes. Its body colour is usually orange, dark red or at least dark



brown-yellow. They are frequently active on soil ground, so it is not unusual to catch them using the pitfall traps. Their forcipulae (appendages of the first trunk segment, used for catching a prey) are wide, with well-developed basal denticle on its claw. The species *S. pusilla* is characterised by low number of pairs of legs and by the morphology of the last pleuropretergite, which is divided, i.e. its intercalary pleurites are not joined with ultimate pretergite, as usual in other species.



Fig. 2: Dorsal site of posterior part of body of male of *Strigamia pusilla*. Arrows indicate division of last pleuropretergite. Orig. László Dányi.

Obr. 2: Hřebetní strana zadního konce těla samce zemivky *Strigamia pusilla*. Šípky označují rozdělený poslední pleuropretergit. Orig. László Dányi.

## IDENTIFICATION KEY FOR THE STRIGAMIA SPECIES REPORTED FROM THE CZECH REPUBLIC

- 1A Ultimate pleuropretergite entire, number of pairs of legs 37 and more ..... 2
- 1B Ultimate pleuropretergite divided into three parts (Fig. 2), number of pairs of legs up to 35...  
..... *S. pusilla*  
*S. pusilla*; maximum body length 24 mm, pairs of legs 33–35
  
- 2A Number of pairs of legs 45 and more ..... 3
- 2B Number of pairs of legs 37–43 ..... *S. acuminata*  
*S. acuminata*; maximum body length 40 mm, pairs of legs 37–43
  
- 3A Anterior sterna with well-developed longitudinal fissure in the middle, number of coxal pores 12–30 ..... *S. crassipes*  
*S. crassipes*; maximum body length 56 mm, pairs of legs 45–59
- 3B Anterior sterna with discrete shallow longitudinal cavity, number of coxal pores 4–9 ..... *S. transsilvanica*  
*S. transsilvanica*; maximum length body 38 mm, pairs of legs 45–57

## ACKNOWLEDGEMENTS

The findings were carried out as a part of malacological survey of the Mazák National Nature Reserve, funded by the OPE project “Implementation of the Natura 2000”, organised and supported by the Nature Conservation Agency of the Czech Republic. We are grateful to Dr. Karel Tajovský (Institute of Soil Biology BC CAS, České Budějovice) for his comments improving quality of manuscript and to Dr. László Dányi (Hungarian Natural History Museum, Budapest) for kind supply of picture of *S. pusilla*.

## REFERENCES

- BONATO L., DÁNYI L., SOCCI A.A. & MINELLI A. (2012): Species diversity of *Strigamia* Gray, 1843 (Chilopoda: Linotaeniidae): a preliminary synthesis. Zootaxa 3593: 1–39.
- BONATO L., EDGECOMBE G.D., LEWIS J.G.E., MINELLI A., PE-REIRA L.A., SHELLEY R.M. & ZAPPAROLI M. (2010): A common terminology for the external anatomy of centipedes (Chilopoda). ZooKeys 69: 17–51.
- BONATO L. & MINELLI A. (2013): Chilopoda Geophilomorpha of Europe: a revised list of species, with taxonomic and nomenclatorial notes. Zootaxa 3770: 1–136.
- DÁNYI L. (2006): Contribution to the Chilopoda fauna of the Maramureş (Romania). Studia Universitatis „Vasile Goldis“, Seria Stiințele Vietii 17: 43–46.
- TUF I.H. & LAŠKA V. (2005): Present knowledge on centipedes in the Czech Republic: a zoogeographic analysis and bibliography 1820–2003. Peckiana 4: 143–161.
- TUF I.H. & TUFOVÁ J. (2008): Proposal of ecological classification of centipede, millipede and terrestrial isopod faunas for evaluation of habitat quality in Czech Republic. Časopis Slezského Muzea v Opavě (A) 57: 37–44.



# OBSAH | CONTENTS

## Články | Articles

- 3 • JANÍČKOVÁ Kateřina & DOLNÍČEK Zdeněk: Šlichová prospekce na lokalitách spojených s údajnou historickou těžbou drahých kovů v Moravskoslezských Beskydech
- 17 • DÍTĚ Daniel, MURIN Bohumil, HÁJEK Michal, HÁJKOVÁ Petra, SEDLÁKOVÁ Blažena, MELEČKOVÁ Zuzana & ELIÁŠ Pavol jun.: Nová lokalita páperca alpínskeho (*Trichophorum alpinum*) na Slovensku
- 23 • DÍTĚ Daniel, MELEČKOVÁ Zuzana & ELIÁŠ Pavol jun.: Ostrica blšná (*Carex pulicaris*) – nový druh vo Veľkej Fatre
- 28 • DŘEVOJAN Pavel & GALUŠKOVÁ Hana: Pozoruhodné floristické nálezy z nivy Moravy mezi Strážnicí a Kunovicemi
- 50 • PLEVOVÁ Lenka: Flóra Vlárského průsmyku v severní části Bílých Karpat
- 77 • ŠTRBA Peter: Poznámky k výškovému rozšíreniu zástupcov radu Polygonales v Západných Karpatoch
- 84 • VALACHOVIČ Milan: Lesné spoločenstvá s borovicou lesnou na Slovensku – koncept ich klasifikácie
- 93 • BANAŠ Marek, STANOVSKÝ Jiří, DVOŘÁK Václav & ZEIDLER Miroslav: Příspěvek k poznání vegetace a fauny brouků 10 let po provedené revitalizaci části potoka Kněžně (k. ú. Prostřední Bečva) v CHKO Beskydy
- 108 • TUŠ Ivan Hadrián & KUPKA Jiří: První nález zemivky *Strigamia pusilla* z území České republiky (Chilopoda: Geophilomorpha)
- 111 • KONVIČKA Ondřej & ČÍŽEK Lukáš: Rozšíření rýhovců *Rhysodes sulcatus* (Fabricius, 1787) a *Omoglymmius germari* (Ganglbauer, 1892) (Coleoptera: Rhysodidae) v České republice
- 115 • GABRIŠ Radim & VÁVRA Jiří Ch.: Nové nálezy *Cerophytum elateroides* (Latreille, 1804) (Coleoptera, Cerophytidae) na severní Moravě s poznámkami k jeho bionomii
- 119 • KONVIČKA Ondřej & SPITZER Lukáš: Zajímavé nálezy brouků (Coleoptera) z východní Moravy ve sbírce Miloslava Herrmanna (Česká republika)
- 123 • TRÁVNÍČEK Dušan: Rozšíření vodomila *Lacobius (Dimorpholaccobius) simulatrix* Orchymont, 1932 v České republice a poznámky k určování tohoto druhu
- 126 • ŠNAJDARA Pavel & SPITZER Lukáš: Rozšíření a stav vybraných populací modráška hořcového Rebelova (*Phengaris alcon rebeli*) (Lepidoptera: Lycaenidae) ve Zlínském kraji (východní Morava)
- 139 • ČELECHOVSKÝ Alois, CHMELA Josef & CHMELA Miloslav: Makrolepidoptera Pozděchova na Valašsku (Česká republika)

## Aktuality a Personálie | Currents News and Personals

- 158 • Moravskoslezská pobočka České botanické společnosti v roce 2015
- 161 • Sítové mapování cévnatých rostlin v okrese Vsetín mimo CHKO Beskydy v roce 2015
- 163 • Vzděláváme multimediálně, interaktivně a efektivně
- 165 • Acta Carpathica Occidentalis. Pokyny pro autory

© Muzeum regionu Valašsko, příspěvková organizace, Vsetín

a Muzeum jihovýchodní Moravy ve Zlíně, příspěvková organizace, 2015

ISBN 978-80-87614-42-6 (Muzeum regionu Valašsko, příspěvková organizace. Vsetín)

ISBN 978-80-87130-39-1 (Muzeum jihovýchodní Moravy ve Zlíně, příspěvková organizace. Zlín)

ISSN 1804-2732